

# BTec National Extended Certificate in Applied Science Summer Independent Learning Y11-12

# Part 1 – Compulsory Content (pages 2-15)

- 1. Watch the videos and complete the notes you may consider adding flashcards / condensed notes, so you can use them to test yourself (metacognition)
- 2. Complete the follow up application questions
- Correct and improve the questions where available (mark scheme at the end of the document)

This will be assessed in the initial assessment

# Part 2 - Highly Recommended (pages 16-18)

This will be beneficial to you and help you succeed in your lessons.

Mark schemes (pages 19-31)

## **Compulsory content**

# **Unit 1: Principles and Applications of Science I**

Answer all the questions. There are links to websites which you may find helpful. You will be given a test on these concepts at the start of the term.

This unit covers some of the key science concepts in biology, chemistry and physics.


This section looks at some of the chemistry concepts you have covered at GCSE and will cover in more depth in Unit 1.

#### Periodicity and properties of elements

#### □ Atomic Structure

https://www.bbc.co.uk/bitesize/guides/zwn8b82/revision/3 (pages 3,4 and 5) https://phet.colorado.edu/sims/html/build-an-atom/latest/build-an-atom\_en.html

#### Q1. Figure 1 shows an atom of element G.



Draw a ring around the correct answer to complete each

- (a) Label A shows
  - an electron an ion a nucleus (1)
- (b) Label B shows
  - an isotope a molecule a neutron (1)
- (c) The atomic number of element G is
- 10 11 (1)
- (d) The mass number of element G is

#### □ Periodic Table


https://www.bbc.co.uk/bitesize/guides/ztv797h/revision/2 (pages 2-8) https://www.rsc.org/periodic-table/

#### **Q2.** The Periodic table below contains **six** errors. Highlight these.

					Н												He
Li	Be											В	С	Ν	0	FI	Ne
nα	Mg											Al	Si	Р	S	CL	Ar
K	Ca	Sc	Ti	٧	Cr	Mn	fe	со	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	У	Zr	Nb	Мо	Тс	Ru	Rh	рD	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ва	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	TI	Pb	Bi	Ро	At	Rn
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg							

(6)

**Q3.** Complete the labels on the diagram below using the following terms:




**Q4.** Read the information below on element **X** carefully. Use this to help you answer the questions which follow.

Element **X** has two different isotopes, both of which contain 17 protons. The least abundant isotope contains 20 neutrons. The second isotope is three time more abundant and contains 2 more neutrons. All the atoms contain 2 electrons in the first shell, 8 electrons in the second shell and 7 electrons in the third.

(a	) Where in the	Periodic	Table is	element <b>X</b>	found:	:
----	----------------	----------	----------	------------------	--------	---

Group: ...... Period: ..... (2)

(b) Use the Periodic Table in Q3. the key and your answer to Q4.(a) to complete Figure 2. for element X



(c) Is element X a metal or non-metal? ......

(4)

(d) Identify an element, in the same group as **X**, which has a lower boiling point than **X**.

......(1)


## □ Chemical reactions and equations

https://www.bbc.co.uk/bitesize/guides/zy4pmsg/revision/1 (pages 1-6) https://www.bbc.co.uk/bitesize/guides/z2bfxfr/revision/1 (pages 1,2)


Equations are used to show chemical reactions.

Reactants are written on the left of the arrow and products are written on the right.

For example:



Atoms cannot be created or destroyed. They are simply rearranged. Therefore, the equation with formulae needs balancing. (You can only add more of the same molecules. You cannot change the formula of any.)



The relative formula mass of a molecule/compound  $(M_r)$  can be calculated by adding the  $A_r$  of all the atoms it contains. The  $A_r$  value for all elements can be found in the Periodic Table.

A<sub>r</sub> of C is 12.0, A<sub>r</sub> of H is 1.0 and A<sub>r</sub> of O is 16.0

$$M_r$$
 of  $C_2H_6 = (2 \times 12.0) + (6 \times 1.0) = 30.0$ 

$$M_r$$
 of  $O_2 = (2 \times 16.0) = 32.0$ 

$$M_r$$
 of  $CO_2 = 12.0 + (2 \times 16.0) = 44.0$ 

$$M_r$$
 of  $H_2O = (2 \times 1.0) + 16.0 = 18.0$ 

(8)

The total mass of the reactants = the total mass of the products

Mass of reactants = 
$$(2 \times M_r C_2 H_6) + (7 \times M_r O_2) = (2 \times 30.0) + (7 \times 32.0) = 284.0$$

Mass of products = 
$$(4 \times M_r CO_2) + (6 \times M_r H_2O) = (4 \times 44.0) + (6 \times 18.0) = 284.0$$

- Q5. Lithium reacts with water to form lithium hydroxide and hydrogen.
  - (a) Balance the symbol equation for this reaction

..... Li(s) + ..... 
$$H_2O(I) \rightarrow$$
 ...... LiOH(aq) +  $H_2(g)$  (1)

(b) (i) Complete the table below for this reaction

	Reactant or product	State	$M_r$
Lithium			
Water	reactant	liquid	18.0
Lithium hydroxide			
Hydrogen			

` '	alculate the ow your v		of the	reacta	ants. A	Are the	ese th	e sam	ne as t	he tot	tal ma	ass c	f the	produ	ucts	?
		 													(2	2)

### Bonding

Chemical reactions involve the breaking and making of bonds. This involves electrons being transferred or shared between atoms.

The total number of electrons at the end of the reaction must be the same as at the start.


**Metal** atoms **lose** electrons and form **positively** charged **ions**.

Non-metal atoms gain electrons and form negatively charged ions

OR by sharing them (in pairs) with another non-metal atom


https://www.bbc.co.uk/bitesize/topics/z33rrwx (ionic compounds, small molecules, metals and alloys)

How do you know which type of bonding is present in an element or compound? Consider the type of element(s) it contains:



**Q6.** The electronic structure of a potassium atom is 2,8,8,1 Draw a diagram to show the electronic structure of a potassium ion. Show the charge on the ion.


**Q7.** Complete the dot and cross diagram to show the electrons in the outer shells of ammonia, NH<sub>3</sub>. Use the periodic table to help you.



(2)

(2)

**Q8.** The diagrams shown an atom of magnesium and an atom of chlorine.



Describe, in terms of electrons, how magnesium atoms and chlorine atoms change into ions to produce Magnesium chloride, MgCl<sub>2</sub>.

You may draw labelled diagrams.

 (4)

# **Unit 2: Practical Scientific Procedures and Techniques**

In this unit you will be required to complete a lot of practical procedures and so it is important that you know about laboratory safety.

## □ Laboratory Safety

 Watch the video on safety in the laboratory: <a href="https://www.youtube.com/watch?v=RhIOYhOvCsQ">https://www.youtube.com/watch?v=RhIOYhOvCsQ</a>

Use this to complete a list of safety rules to follow when completing any experiment.

3.	
7.	
8.	


















(8)


You will be using a number of different chemicals and apparatus when completing these experiments.

- Follow the instructions provided to complete the table below on hazard symbols
  - Match the old hazard symbol to the new symbol.
  - ii) Match the new hazard symbol to the hazard name.

    <a href="https://www.acs.org/content/dam/acsorg/about/governance/committees/chemicalsafety/publications/acs-secondary-safety-guidelines.pdf">https://www.acs.org/content/dam/acsorg/about/governance/committees/chemicalsafety/publications/acs-secondary-safety-guidelines.pdf</a> (page 22 and 23)</a>
  - List the precautions which should be taken (in addition to wearing a labcoat and safety glasses) when handling chemicals with these hazards to minimise the chance of an accident occurring.

    <a href="https://www.acs.org/content/dam/acsorg/about/governance/committees/chemicalsafety/publications/acs-secondary-safety-guidelines.pdf">https://www.acs.org/content/dam/acsorg/about/governance/committees/chemicalsafety/publications/acs-secondary-safety-guidelines.pdf</a> (pages 38-40)





Name	Precautions
harmful / irritant	
oxidising agent	
flammable	
harmful to the environment	
corrosive	
toxic	
	(19)

## □ Practical techniques

One of the practical techniques you will need to complete is the preparation of a standard solution and performing a titration to test the solution you have prepared.


 Watch these videos to help you answer the questions <a href="https://www.youtube.com/watch?v=xBKyjXUhJy0">https://www.youtube.com/watch?v=xBKyjXUhJy0</a>
 <a href="https://www.youtube.com/watch?v=gzvzvDv\_BnA">https://www.youtube.com/watch?v=gzvzvDv\_BnA</a>

Q1. (a)	What is a standard	d solution? .	 	 	 


(b) The diagram below shows the apparatus used to make a standard solution of sodium carbonate.

Complete the labels. (7)

.....(1)



(c) The standard solution prepared can be used to find the concentration of a solution of hydrochloric acid.



- (i) Complete the label to show name of the apparatus in which the acid is placed. (1)
- (ii) What is the name given to this procedure? ......(1)
- (iii) Figure 2. shows the level of the sodium carbonate solution in the burette at the start and the end of one titration. Use these to work out the volume of sodium carbonate added in the titration. Give your answer to 2 d.p.


Volume 
$$Na_2CO_3(aq)$$
 added = ..... cm<sup>3</sup> (1)

Another practical technique you will need to complete is chromatography.

 The links below may help you to answer the questions on this technique. <a href="https://www.youtube.com/watch?v=lj5OWzhZSac">https://www.youtube.com/watch?v=lj5OWzhZSac</a>
 <a href="https://www.bbc.co.uk/bitesize/quides/z9dfxfr/revision/4">https://www.bbc.co.uk/bitesize/quides/z9dfxfr/revision/4</a>

Q2.	<b>(a)</b> What	is chro	matogr	aphy us	ed for?	 	 	 	 	
						 	 	 	 	(1)


(b) A student used paper chromatography to analyse a black food colouring. They placed spots of known food colours, A, B, C, D and E and the black food colouring on a sheet of chromatography paper. They set up the apparatus as shown in **Diagram 1**.



The student made **two** errors in setting up the apparatus. Identify the **two** errors and describe the problem each error would cause.

| <br> | <br>. (4) |
|------|------|------|------|------|------|------|------|------|------|-----------|

(c) A different student set up the apparatus without making any errors. The chromatogram in **Diagram** 2. shows the student's results.



(i) What do the results tell you about the composition of the black food colouring?

/ii\	Use Diagram 2. to complete Table 1.
(11)	USE Diagram 2. to complete rable 1.

Table 1.

	Distance in mm
Distance from start line to solvent front	
Distance moved by food colour C	

	(iii)	) Use your answers in (c)(ii) to calculate the Rf value for food colour C. Sh	low your working	าตร
--	-------	---	------------------	-----

R<sub>f</sub> value = ..... (1)

(2)

(iv) Table 2. gives the results of chromatography experiments that were carried out on some known food colours, using the same solvent as the students.

Table 2.

Name of food colour	Distance from start line to solvent front in mm	Distance moved by food colour in mm	R <sub>f</sub> value
Ponceau 4R	62	59	0.95
Carmoisine	74	45	0.61
Fast red	67	27	0.40
Erythrosine	58	17	0.29

hich of the food colours in <b>Table 2.</b> could be food colour <b>C</b> from the chromatogram? Give the ason for your answer.	ne
	. (2)

	Obtaining a	nd analysing	results	obtained in	an experimen
_	Obtaining a	ına anaryəniç	j i Couito	obtained in	an experimen

It is important to keep a record of all data whilst carrying out practical work. It is good practice to draw a table before starting the experiment and then enter results straight into the table.

Tables should have clear headings with units.

Time / min	Temperature / °C
0	27.6
1	27.4
2	27.2

The independent variable is the left-hand column in a table, with the following columns showing the dependent variables. All measurements should be written to the same number of decimal places (matching the precision of the measuring instrument).

https://www.bbc.co.uk/bitesize/guides/zcxp6yc/revision/1https://www.bbc.co.uk/bitesize/guides/zcxp6yc/revision/6

**Q3.** A student was told to complete a practical to investigate how temperature affects the rate of a reaction. The student carried out the reaction at five different temperatures and recorded the time taken for each.

The student then calculated the rate of reaction, in s<sup>-1</sup> for each experiment using the equation:

The student's results and calculations are shown below:

at 24.5 °C the experiment took 340 seconds	1/340 = 0.0029 s
at 39.0 °C ít took 256 sec	1/256 = 0.0039 s
at 58.0 °C the experiment took 124 s	$1/124 = 0.0081  s^{-1}$
80.5 °C 62 s	1/62 = 0.0161
51 °C 186 s	1/186 = 0.0054

(a) What is the independent variable in this experiment? Circle the correct answer

rate of reaction time temperature (1)

(4)


**(b)** Tabulate the student's data in an appropriate manner.

#### https://www.bbc.co.uk/bitesize/guides/z8fq6yc/revision/8

Drawing a graph of the results obtained usually makes it easier to interpret the data and draw conclusions.

The independent variable is shown on the *x*-axis and the dependent variable is shown on the *y*-axis.


Axes should always be labelled with the quantity being measured and the units.



Data points should be marked with a cross, x.

When choosing the scales consider:

- the maximum and minimum values of each variable.
- whether 0,0 should be included as a data point.
- how to draw the axes without using difficult scale markings (e.g. multiples of 3, 7, etc)
- the data points should cover at **least half** of the grid supplied for the graph.



Consider the following when deciding where to draw a line of best fit:

- the line can be straight or curved
- the line should pass through, or very close to, the majority of plotted points (ignoring any anomalous points)
- for points not on the line make sure that there are as many points on one side of the line as the other
- the line should be continuous and drawn with a sharp pencil (use a rule for a straight line)
- the line will go through the origin (0,0) if a value of 0 for the independent variable would produce a value of 0 for the dependent variable

**Q4.** A student investigated how the temperature of a metal block changed with time.

An electric heater was used to increase the temperature of the block.

The heater was place in a hole drilled in the block as shown in Figure 1.

Power supply

Electric heater

Metal block

Heat-resistant mat

The student measured the temperature of the metal block every 60 seconds. **Table 3.** shows the student's results.

Table 3.


Time in s	Temperature in °C
0	20.0
60	24.5
120	29.0
180	31.0
240	31.5

(a) Complete the graph of the data from Table 3. on Figure 2.

- Choose a suitable scale for the x-axis.
- Label the x-axis and label the y-axis.
- Plot the student's results.
- Draw a line of best fit.

(5)

Figure 2.



**(b)** Use the graph to find the temperature of the metal block at time 100 s.

Temperature at 100s = .....°C (1)

;	Determine the gradient of the graph over the first 60 seconds.
	Gradient =°C / s (2)

PTO for highly recommended content

# **Highly recommended content**

Make notes from the following resources, then have a go at completing the questions

## Titrations and mole calculations

https://www.bbc.co.uk/bitesize/guides/zx98pbk/revision/3

https://www.youtube.com/watch?v=wPGVQu3UXpw







https://www.youtube.com/watch?v=ovx-Sro4NXM

**Q1.**This question is about acids and alkalis.

(a) Dilute hydrochloric	acid is a strong acid.
-------------------------	------------------------

Explain why an acid can be described as both strong	g and dilute.

(2)

A student titrated 25.0 cm³ portions of dilute sulfuric acid with a 0.105 mol/dm³ sodium hydroxide solution.

(c) The table below shows the student's results.

	Titration	Titration	Titration	Titration	Titration
	1	2	3	4	5
Volume of sodium hydroxide solution in cm <sup>3</sup>	23.50	21.10	22.10	22.15	22.15

The equation for the reaction is:

$$2 \text{ NaOH} + \text{H}_2\text{SO}_4 \rightarrow \text{Na}_2\text{SO}_4 + 2 \text{ H}_2\text{O}$$

Calculate the concentration of the sulfuric acid in mol/dm<sup>3</sup>

- <u></u>		
	Concentration of sulfuric acid =	mol/dm <sup>3</sup>
	tudent should use a pipette to measure the dilute sulfuric acid a	
	tudent should use a pipette to measure the dilute sulfuric acid a	
	tudent should use a pipette to measure the dilute sulfuric acid a	
	tudent should use a pipette to measure the dilute sulfuric acid a	
burette to measur	tudent should use a pipette to measure the dilute sulfuric acid a	
Calculate the mas	tudent should use a pipette to measure the dilute sulfuric acid a re the sodium hydroxide solution.	
Calculate the mas	tudent should use a pipette to measure the dilute sulfuric acid a re the sodium hydroxide solution.  ss of sodium hydroxide in 30.0 cm³ of a 0.105 mol/dm³ solution.	
Calculate the mas	tudent should use a pipette to measure the dilute sulfuric acid a re the sodium hydroxide solution.  ss of sodium hydroxide in 30.0 cm³ of a 0.105 mol/dm³ solution.	
Calculate the mas	tudent should use a pipette to measure the dilute sulfuric acid a re the sodium hydroxide solution.  ss of sodium hydroxide in 30.0 cm³ of a 0.105 mol/dm³ solution.	
Calculate the mas	tudent should use a pipette to measure the dilute sulfuric acid a re the sodium hydroxide solution.  ss of sodium hydroxide in 30.0 cm³ of a 0.105 mol/dm³ solution.	

Use only the student's concordant results.

Describe how a sample of copper chloride crystals could be made from copper carbonat and dilute hydrochloric acid.	
A student wanted to make 11.0 g of copper chloride.	
The equation for the reaction is:	
$CuCO_3 + 2HCI \rightarrow CuCl_2 + H_2O + CO_2$	
Relative atomic masses, A <sub>r</sub> : H = 1; C = 12; O = 16; Cl = 35.5; Cu = 63.5	
Calculate the mass of copper carbonate the student should react with dilute hydrochloric acid to make 11.0 g of copper chloride.	2
Mass of copper carbonate =	_ (
The percentage yield of copper chloride was 79.1 %.	
Calculate the mass of copper chloride the student actually produced.	
Actual mass of copper chloride produced =	,

**Q2.** A student investigated the reactions of copper carbonate and copper oxide with dilute hydrochloric acid.

#### Mark Scheme

### Periodicity and properties of elements

#### ☐ Atomic Structure

https://www.bbc.co.uk/bitesize/guides/zwn8b82/revision/3 (pages 3,4 and 5) https://phet.colorado.edu/sims/html/build-an-atom/latest/build-an-atom/en.html

Q1. Figure 1 shows an atom of element G.

Figure 1

A

Proton
B

Draw a ring around the correct answer to complete each sentence.

(a) Label A shows

an electron an ion a nucleus (1)
(b) Label B shows

an isotope a molecule a neutron

(c) The atomic number of element G is

(5) 6 10 11 16 = number of protons (1)


(d) The mass number of element G is

5 6 10 (1) 16 5 protous + Eventrous (1)

#### ☐ Periodic Table

https://www.bbc.co.uk/bitesize/guides/ztv797h/revision/2 (pages 2-8) https://www.rsc.org/periodic-table/


Q2. The Periodic table below contains six errors. Highlight these.



CAPITAL their lawer case

(6)

(1)



Fluorine

(1)

(d) Identify an element, in the same group as X, which has a lower boiling point.

Q5. Lithium reacts with water to form lithium hydroxide and hydrogen.

(a) Balance the symbol equation for this reaction

..2. 
$$Li(s) + ...2. H_2O(1) \rightarrow ...2.. LiOH(aq) + H_2(g)$$
 (1)


(b) (i) Complete the table below for this reaction

	Reactant or product	State	Mr
Lithium	reactant	said	
Water	reactant	liquid	18.0
Lithium hydroxide	product	agreeus (soluti	an) 23.9
Hydrogen	product	903	20


(ii) Calculate the total mass of the reactants. Are these the same as the total mass of the products? Show your workings.


Q6. The electronic structure of a potassium atom is 2,8,8,1


Draw a diagram to show the electronic structure of a potassium ion. Show the charge on the ion.



Q7. Complete the dot and cross diagram to show the electrons in the outer shells of ammonia, NH<sub>3</sub>. Use the periodic table to help you.








(2)


(8)

Q8. The diagrams shown an atom of magnesium and an atom of chlorine.



Describe, in terms of electrons, how magnesium atoms and chlorine atoms change into ions to produce Magnesium chloride, MgCl<sub>2</sub>.

You may draw labelled diagrams.



One Mg atom loses Le to one Cl atam
It loses a 2nd e to a 2nd e atom
Mg forms the Mg2+ ian and each CI forms a Ct ian

## Unit 2: Practical Scientific Procedures and Techniques


In this unit you will be required to complete a lot of practical procedures and so it is important that you know about laboratory safety.

#### □ Laboratory Safety

 Watch the video on safety in the laboratory: https://www.youtube.com/watch?v=RhIOYhOvCsQ

Use this to complete a list of safety rules to follow when completing any experiment.

1. Wear a lab cost (buttaned up)
2. Wear sixehy glamen / giggles
3. The back lang hair (pasticulally when using a Bunsen burner)
4. White a vish anemment (begane your complete the practical)
5. Keep the lab tidy.
6. Wipe up spillages
7. Do not eat (or drink or chemique) in the lab
8. Wash you hand (pasticularly begane you leave the lab)




## Practical techniques

One of the practical techniques you will need to complete is the preparation of a standard solution and performing a titration to test the solution you have prepared.


 Watch these videos to help you answer the questions https://www.youtube.com/watch?v=xBKyjXUhJy0 https://www.youtube.com/watch?v=rLc148UCT2w https://www.youtube.com/watch?v=gzvzvDv BnA

Q1. (a) What is a standard solution? ... It is a solution of (accurately) known.

(b) The diagram below shows the apparatus used to make a standard solution of sodium carbonate. Complete the labels.



(c) The standard solution prepared can be used to find the concentration of a solution of hydrochloric acid.




- (i) Complete the label to show name of the apparatus in which the acid is placed. (1)
- (iii) Figure 2. shows the level of the sodium carbonate solution in the burette at the start and the end of one titration. Use these to work out the volume of sodium carbonate added in the titration. Give your answer to 2 d.p.

Volume Na<sub>2</sub>CO<sub>3</sub>(aq) added = ....22..\5....... cm<sup>3</sup> (1)

8

Q2. (a) What is chromatography used for? ... to separate fand analyse) components ...

(b) A student used paper chromatography to analyse a black food colouring. They placed spots of known food colours, A, B, C, D and E and the black food colouring on a sheet of chromatography paper. They set up the apparatus as shown in Diagram 1.



The student made **two** errors in setting up the apparatus. Identify the **two** errors and describe the problem each error would cause.

Stat hive drawn in inte

so it will run I dissolve in the solvent

(1)


Solvent above the stat line / spots under the solvent

so they will worsh aff the paper/ mix with the solvent

(1)

(c) A different student set up the apparatus without making any errors. The chromatogram in **Diagram** 2. shows the student's results.

Diagram 2



(i) What do the results tell you about the composition of the black food colouring?

The black food charing contains A and E (1)
and are other (unknown) substance (1) (2)

(ii) Use Diagram 2. to complete Table 1.

(2)

Table 1.

	Distance in mm
Distance from start line to solvent front	28.5 / 29
Distance moved by food colour C	allow 11-12

(iii) Use your answers in (c)(ii) to calculate the R<sub>f</sub> value for food colour C. Show your workings.

$$R_f$$
 value = .0.38 - 0.42...(1)

(iv) Table 2. gives the results of chromatography experiments that were carried out on some known food colours, using the same solvent as the students.

Table 2.

Name of food colour	Distance from start line to solvent front in mm	Distance moved by food colour in mm	R <sub>f</sub> value
Ponceau 4R	62	59	0.95
Carmoisine	74	45	0.61
Fast red	67	27	0.40
Erythrosine	58	17	0.29

Which of the food colours in **Table 2**, could be food colour **C** from the chromatogram? Give the reason for your answer.

	allas nave
	b.) if he not cause as any
/ /	

Q3. A student was told to complete a practical to investigate how temperature affects the rate of a reaction. The student carried out the reaction at five different temperatures and recorded the time taken for each.

The student then calculated the rate of reaction, in s<sup>-1</sup> for each experiment using the equation: rate of reaction = 1


The student's results and calculations are shown below:

at 24.5 °C the experiment took 340 seconds	1/340 = 0.0029 s-1
at 39.0 °C it took 256 sec	1/256 = 0.0039 s-1
at 58.0 °C the experiment took 124 s	1/124 = 0.0081 s-1
80.5 °C 62 s	1/62 = 0.0161
51 °C 186 s	1/186 == 0.0054

(a) What is the independent variable in this experiment? Circle the correct answer

	rate of reaction	time	(temperature)	(1)
(b) Tabulate the student's data in an appropriate manner.			ner.	(4)

temperature 1°C	time1s.	rate of reaction 1.5.
24,5	340	0.0029
39.0	256	0.0039
51.0	186	0.0054
58.0	124	0,0081
80.5	62	0.0161



	The rate of change of temperature of the block is given by the gradient of the graph.  Determine the gradient of the graph over the first 60 seconds.
	24.(-0) = 0.41(1)
**	60-0 (1)
	Gradient =0.4)°C / s (2)

write to ldp to world

## **Highly recommended content**

# Q1. (strong because) completely ionised (in aqueous solution) (a) ignore pH allow dissociated for ionised do not accept hydrogen is ionising do not accept H+ are ionised 1 (dilute because) small amount of acid per unit volume ignore low concentration 1 (c) (titre): chooses titrations 3, 4, 5 1 average titre = 22.13 (cm<sup>3</sup>) allow average titre = 22.13(3...) (cm<sup>3</sup>) allow a correctly calculated average from an incorrect choice of titrations 1 (calculation): (moles NaOH = $\frac{22.13}{1000} \times 0.105 = 0.002324$ allow use of incorrect average titre from step 2 1 (moles $H_2SO_4 =$ $\frac{1}{2} \times 0.002324 = 0.001162$ allow use of incorrect number of moles from step 3 1 (concentration = 0.<u>001162</u>×1000) $= 0.0465 \text{ (mol/dm}^3)$ allow use of incorrect number of moles from step 4 1 alternative approach for step 3, step 4 and step 5 $\frac{2}{1} = \frac{22.13 \times 0.105}{25.0 \times conc. H_2 SO_4} (1)$ (concentration $H_2SO_4 =$ ) 22.13 × 0.105 25.0 × 2 $= 0.0465 \, (mol/dm^3) \, (1)$ an answer of 0.046473 or 0.04648 correctly rounded to at least 2 sig figs scores marking points 3, 4 and 5 an answer of 0.092946 or 0.09296 or 0.185892 or

0.18592 correctly rounded to at least 2 sig figs scores

marking points 3 and 5 an incorrect answer for one step does **not** prevent allocation of marks for subsequent steps

(d) pipette measures a fixed volume (accurately)

1

(but) burette measures variable volume allow can measure drop by drop

1

(moles =)  $\frac{30}{1000} \times 0.105$ or 0.00315 (mol)

(mass per dm<sup>3</sup> =)  $0.105 \times 40$ or 4.2 (g)

1

$$(mass = \frac{30}{1000} \times 0.105 \times 40)$$

1

= 0.126 (g)

an answer of 0.126 (g) scores **2** marks an answer of 126(g) scores **1** mark an incorrect answer for one step does **not** prevent allocation of marks for subsequent steps

[12]

**Q2.** 

(a) add excess copper carbonate (to dilute hydrochloric acid)

accept alternatives to excess, such as 'until no more reacts'

1

filter (to remove excess copper carbonate)

reject heat until dry

1

heat filtrate to evaporate some water **or** heat to point of crystallisation accept leave to evaporate or leave in evaporating basin

1

leave to cool (so crystals form)
until crystals form

1

must be in correct order to gain 4 marks

(b)  $M_r \text{ CuCl}_2 = 134.5$ 

correct answer scores 4 marks

1

moles copper chloride = (mass /  $M_r$  = 11 / 134.5) = 0.0817843866

1

	<i>M</i> <sub>r</sub> CuCO₃= 123.5	1
	Mass CuCO <sub>3</sub> (=moles × M <sub>2</sub> = 0.08178 × 123.5) = 10.1(00)	1
	accept 10.1 with no working shown for 4 marks	•
(c)	$\frac{79.1}{100} \times 11.0$	
	or	
	$11.0 \times 0.791$	1
	8.70 (g)	1

accept 8.70(g) with no working shown for 2 marks